

R008-09

C 会場 : 11/26 PM2 (14:50-16:20)

15:20~15:35:00

#松清 修一¹⁾, 諫山 翔伍¹⁾, 佐藤 弓真¹⁾, 高橋 佳夏¹⁾, 森田 太智¹⁾, 竹崎 太智²⁾, 山崎 了³⁾, 田中 周太³⁾, 佐野 孝好⁴⁾, 蔵満 康浩⁴⁾, 境 健太郎⁵⁾, 坂和 洋一⁴⁾

(¹⁾ 九大, (²⁾ 富山大, (³⁾ 青学大, (⁴⁾ 阪大, (⁵⁾ 核融合研

Reflected ions and nonstationarity confirmed in collisionless shock experiment using power laser

#Shuichi Matsukiyo¹⁾, Shogo Isayama¹⁾, Yuma Sato¹⁾, Kana Takahashi¹⁾, Taichi Morita¹⁾, Taichi Takezaki²⁾, Ryo Yamazaki³⁾, J. Shuta Tanaka³⁾, Takayoshi Sano⁴⁾, Yasuhiro Kuramitsu⁴⁾, Kentaro Sakai⁵⁾, Youichi Sakawa⁴⁾

(¹⁾Kyushu University, (²)University of Toyama, (³)Aoyama Gakuin University, (⁴)Osaka University, (⁵)National Institute for Fusion Science

We reproduce a collisionless shock using high-power laser experiment which is increasingly being recognized as a major empirical research tool in this field. Long pulse laser irradiates a plate target to create a high-speed plasma flow normal to the target plate. Surrounding gas is ionized with intense radiation generated from the interaction between the laser and the target material. Ambient magnetic field almost parallel to the plate surface is applied by using a Helmholtz-like coil so that the gas plasma was nearly homogeneously magnetized. The high-speed target plasma acts as a magnetic piston to form a shock in the ambient gas plasma.

By analyzing the spatiotemporal imaging data of self-emission intensity, we extracted fluctuations of the shock front and demonstrated that their period is on the order of the inverse of the ion gyro frequency. Thomson scattering measurement in the transition region of the shock showed highly asymmetric spectrum, indicating that the local plasma is composed of two ion populations having different bulk speed.

We developed 2D PIC simulation to reproduce collective Thomson scattering in a beam-plasma system in a self-consistent manner. Characteristics of 2D spectrum of the scattered waves for various beam parameters were examined. Assuming weak beam-plasma condition, the reproduced scattered wave spectrum showed asymmetric nature similar to what was observed in the experiment. Therefore, we conclude that the local plasma in the shock transition region observed in the experiment is composed of background ions, electrons, and beam (or reflected) ions. The presence of the weak reflected ion beam may cause the fluctuations of the shock front. The fluctuations may be attributed to breathing, a phenomenon observed in collisionless shocks with high ion temperatures.